

# **Spatial distribution of the near coast and onshore seismicity of Crete (South Greece) with special emphasis to Heraklion basin (Central Crete)**

KOKINO ELEN<sup>a</sup>, VALLIANATOS PHILIPPOS<sup>a</sup>, SARRIS APOSTOLOS<sup>b</sup>, MOISIDI MARGARITA<sup>a</sup>, TZANAKI IFIGENEIA<sup>a</sup>, TZAKALAKI EFTIMIA<sup>a</sup>, TZISKAKI EVANGELIA<sup>a</sup>

<sup>a</sup>Laboratory of Geophysics and Seismology, Department of Natural Resources and Environment, Technological Educational Institute Crete, 3 Romanou Str. Chalepa, Chania, Crete, GR 73133, GREECE, ekokinou@chania.teicrete.gr

<sup>b</sup>Laboratory of Geophysical-Satellite Remote Sensing & Archaeo-environment, Institute for Mediterranean Studies, Foundation for Research & Technology - Hellas, P.O. Box 119, Rethymnon 74100, Crete, Greece, asaris@ret.forthnet.gr

*Abstract:* - In the context of the present work the spatial distribution of the seismic activity around Crete Island in the Southern Hellenic Arc for the period 2003-2007 is studied. Especially in the period July-October 2007 about 600 events with low and moderate magnitudes have been indicated. Most of these events are located near coast and onshore of Crete. Additionally, we studied the relation of the earthquake epicenters with specific fault zones for the Heraklion basin.

GIS techniques were used for mapping the distribution of earthquake epicenters on the various topographic and geological features of the area. Maps were created through interpolation algorithms. Spatial tools and statistical analysis were exploited to examine the correlation between earthquake loci and faulting orientations. The distribution of the epicenters indicates that the near coast and onshore seismicity is closely associated to the Crete tectonics. Especially for Heraklion prefecture the epicenter distribution is related to E-W, NE-SW and NW-SE striking faults.

*Key-Words:* - Earthquake, geotectonic structure, normal faulting, GIS


## **1 Introduction**

The Hellenic Arc represents the most seismically active area of Europe due to the interaction between Eurasia and Africa plates. Main geotectonic feature of the area is the Hellenic Trench, where the eastern Mediterranean oceanic lithosphere (frontal part of the African plate) is subducted under the Aegean overriding plate. Earthquakes with magnitudes of up to 8.0 have been reported in the literature since the early historic times [1] [2] pointing out the great seismogenetic potential of the area. According to [3], an ocean-continent interaction occurs on a curved subduction zone, which is characterised by a shallow branch (20.0-100.0 km) of the Wadati-Benioff zone, intersecting the outer side of the sedimentary arc (Western Peloponnesos-west of Kythira-south coast of Crete, east coast of Rhodes) and dips at low angle (~30°) to the north and northeast. Subduction at the Hellenic subduction zone appears to have been operated continuously since the late Cretaceous [4] [5].


The Hellenic Arc can be at present subdivided into three zones with different kinematic behaviour: In

the northwestern part of the Aegean region, the boundary between the Aegean and African plates (e.g Ionian Sea) is of continent-continent type now due to the collision of the Hellenides with the Apulian platform [6] [7] [8]. The boundary between these regions of contrasting subduction style is presently formed by the right-lateral Kefallonia Transform Fault Zone [9]. The southern part of the Hellenic arc south of, and including Crete and Rhodos has since the early Pliocene been associated with left-lateral strike slip along the Pliny and Strabo trenches [10]. In between these two regions, active northeast directed subduction is accommodated along the Hellenic Trench and deformation in the overriding plate on the Peloponnesos, and in the Kythira-Antikythira strait [11] is associated with a complex pattern of arc-parallel and arc-normal extension and strong compression perpendicular to the Hellenic Trench (Fig 1a) [12]. Beneath this area, the subduction zone dips gently to the northeast, changing to a steeper dip farther northeast beneath the Argolikos Gulf [13].

Scope of this work is to study the spatial distribution of the near coast and onshore earthquakes occurred in the period 2003-2007 and to associate the earthquake loci with specific geotectonic features. Innovative GIS techniques were used in order to implement the pre-mentioned scope. The Heraklion basin, as a well studied region, was selected in order to combine the distribution of the near coast and onshore seismicity with the orientation of specific fault zones.



(a)



(b)

Figure 1 The study area (a) and the seismic network (b) of the laboratory of Geophysics and Seismology of the Technological educational Institute of Crete

## 2 Geological setting of Crete

The island of Crete (Fig.1a) represents an emergent high in the fore-arc of the Hellenic Subduction Zone, indicating the transition between African and Eurasian plates. A variety of intensive studies in the last decades figured out the geodynamic attributes of the wide area of Southern Hellenic Arc [6]. Additionally [6] and [14] concluded that the north-dipping Wadati-Benioff seismic zone is extending beneath Crete to a depth of about 200 Km. Nowadays the Hellenic arc is associated with moderate arc-parallel extension and strong compression perpendicular or oblique to it.

According to tomographic studies the seismicity of the wide area around Crete corresponds to a lithospheric slab extending through the transition zone and into the lower mantle below Europe. The Mediterranean Ridge (MR) complex consists of accumulated sediments of the subducted African plate. Between Crete and the MR are located a series of E-NE trending depressions or troughs (e.g. Hellenic, Ptolemy, Pliny, Strabo, Fig. 1a).

North of the Crete Island, the tomography drops off into the thinned continental crust of the Cretan sea [15]. The volcanic arc is positioned about 100 Km north of Crete, represented by the island of Santorini. GPS and seismic studies [16] show that Crete and the South Aegean are moving together as a coherent block. The divergent motion between the Aegean block and mainland Europe is indicated by an extension zone, with Crete and Aegean diverging from mainland Europe at a rate of about 3 cm/yr [17] while Africa is moving northward relative to Europe at a rate of about 1 cm/yr [18].

According to previous geologic and tectonic studies [6] [19] three successive fault groups-generations could be distinguished on Crete Island and especially in Heraklion basin (Fig. 2). The first group represents E-W trending faults, kilometer-scale, mainly cutting the basement rocks or bound basement rocks and Miocene sediments. The second group consists of large and moderate-scale N-S striking faults, cutting the previous mentioned group. The third group comprises Kilometer-scale faults striking NE-SW and NW-SE which appear to be the youngest faults occurring on Crete Island.

The largest earthquakes occurred on and around Crete [3] indicate E-W extension along N-S striking faults onshore Crete. Fault plane solutions in the western part of Crete Island found T-axes nearly horizontal and oriented mostly E-W in the upper crust [20] [21]. [22] studied large to moderate earthquakes along the Hellenic Arc and noticed that fault plane solutions for half of the 15 studied events show approximately E-W compression, sub-parallel to the trend of the eastern Hellenic Trench. N-S compression dominates the offshore regions south of Central and Western Crete, while normal and thrust faulting is the predominant mechanism south of Eastern Crete.




Figure 2 Faults of the Heraklion Prefecture

### 3 Methodology

*Topology of the seismic network* The new telemetric seismic network (Fig. 1b) has been installed since the end of 2003 on the island of Crete and the broader area of South Aegean and is continuously operated by the Laboratory of Geophysics and Seismology of the Technological Educational Institute of Crete in order to provide modern instrumental coverage of seismicity in the southern Greece, as well as some more insight into the stress and deformation fields, tectonics, structure and dynamics of the Hellenic Arc. Network's geometry as well as site selection has been chosen carefully, since the primary goal is to locate seismic events, fact which assures the most accurate determination of seismic parameters. Furthermore, plenty of studies are conducted, concerning mainly the crust structure of the area which will lead to a representative velocity model, the stress-field and the focal depth distribution. The South Aegean Seismic Network (SASN) is now well established with a stable technical and financial position. The associated SASN database has increased in both quantity and quality during the period 2003-2008 and is being extensively used for research. Due to the technical development of the network, there has also been a good knowledge of software applications for seismic data acquisition, communication and processing.

*Earthquake dataset* The earthquakes of magnitude  $M \geq 2.5$  which occurred in the broader area of Crete, during the period 2003-2007 have been selected in the context of the present work. These data are based on the recordings at two permanent seismological networks in Greece (National Observatory of Athens - Institute of Geodynamics

NOA, and the SASN) as well as the Eastern Mediterranean Seismological Centre (EMSC). The stations (Fig. 1b) of SASN that used in the present work are mostly distributed in the periphery of Crete as well the neighboring islands. Furthermore the earthquake epicenters (Fig. 3a), recorded by the SASN in the period 2006-2007, are distributed offshore and onshore Crete. The density map of the earthquake epicenters (Fig. 3b) shows an intense near coast and onshore seismicity that gave rise to study the relation between the spatial distribution of the faults and the earthquake epicenters for the area of the Heraklion prefecture contacting more intense earthquake analyses for two distinct periods, namely July-August 2007 and September-October 2007.




Figure 3 a) Distribution of the earthquakes epicenters monitored by the TEI network for the period 2006-2007, b) density map of the earthquake epicenters

*GIS techniques and mapping* Density maps were created for each one of the NOA, EMSC and TEI seismic activity datasets corresponding to the periods of 2003-2006, and 2006-2007 respectively. A kernel density algorithm using a quadratic formula where the highest value corresponds to the center location falling to zero at the search radius distance was preferred in the calculations of the

density maps for creating smoother distribution surfaces. A circular search area of approximately 10km was used around the location of each seismic epicenter. The resulting density surfaces having a resolution of about 1km indicate the spread of the seismic epicenters over the areas surrounding the island of Crete. Similar density maps were created for the micro-seismicity of the region for two distinct periods, namely July-August 2007 and September-October 2007.

In order to correlate the seismic risk of the prefecture of Heraklion in terms of the activated inland faults of the region, the IGME (Institute for Geological and Mineral Exploration) geological maps of the region were digitized and the described faults were classified in four main categories depending on their alignment, namely E-W, N-E, NW-SE and NE-SW. A buffer distance of 500m was used around each epicenter for the seismic incidences corresponding to the NOA, TEI and micro-seismicity (whole period of July-October 2007), each of which was approached independently. Similarly, buffer zones of 250m were created around each one of the fault classes. Boolean algebra was then used to combine the above surfaces and generate areas that overlapped. In this way, it was possible to correlate the seismic activity of the area with specific faults (or even sections of faults) and examine the attributes of specific seismic zones.

#### 4 Results

Figure 4a shows the spatial distribution of the earthquake epicenters recorded in the period July-October 2007 while Figures 4b and 4c present the density maps of the earthquakes for two distinct periods, namely July-August 2007 and September-October 2007 respectively. In the period July-August 2007 an intense seismicity is observed in western Crete. A N-S orientation of the epicenters is clearly seen in the westernmost part of Crete while an E-W orientation of the epicenters is present at the northeastern part of Crete. The seismicity pattern of the period September-October shows an intense onshore seismicity in the whole island with no specific orientations.

Thereinafter we have chosen the area of Heraklion Prefecture in order to study the relation between fault orientation and epicenters. Figures 5 to 8 present the E-W, N-S, NW-SE and NE-SW activated faults for the period 2003-2007 based on NOA and EMSC recordings. All fault-group generations show to be responsible for the seismicity

of the study area. Additionally seismicity decreases rapidly from east to west with practically no events along the western boundary of Heraklion basin. Especially, E-W orientated faults influence the seismicity of the whole study area while N-S orientated faults show to be responsible for the seismicity in the northern part of Heraklion Prefecture. NW-SE and NE-SW fault groups are probably responsible for the seismicity in the whole study area.




Figure 4 a) Distribution of the earthquake epicenters monitored by the TEI network for the period July-October 2007, (b) Density map of the earthquakes recorded in the period July-August 2007, (c) Density map of the earthquakes recorded in the period September-October 2007

The maps of Figures 9 and 10 present the general seismicity pattern in relation to all fault-group generations for the periods 2003-2007 and July-October 2007 respectively. At this point we must refer that for the period July-October 2007 the recordings of the SASN were used and intensively analyzed. Both maps indicate that E-W, NW-SE and NE-SW fault orientations can generally be considered responsible for the seismicity of the study area.




Figure 5 E-W Activated faults of the Heraklion Prefecture for the period 2003-2007




Figure 6 N-S Activated faults of the Heraklion Prefecture for the period 2003-2007




Figure 7 NW-SE Activated faults of the Heraklion Prefecture for the period 2003-2007




Figure 8 NE-SW Activated faults of the Heraklion Prefecture for the period 2003-2007




Figure 9 Activated faults of the Heraklion Prefecture for the period 2003-2007




Figure 10 Activated faults of the Heraklion Prefecture for the period July-October 2007

## 5 Discussion and Conclusions

In the context of the present work the earthquake data from the catalogs of NOA and EMSC as well as recordings of the SASN network were processed in order to study the near coast and onshore spatial distribution of the seismicity in Crete. Additionally the relation between fault-group generations and epicenters was investigated for the area of Heraklion Prefecture located in central Crete. Significant seismic activity, possibly of low magnitude, characterizes the near coast and onshore part of Crete. The westernmost part of Crete is characterized by an N-S orientation of the earthquake epicenters while in the rest of the island an E-W trend in the orientation of the epicenters is observed. Especially for the area of the Heraklion basin the seismic activity is concentrated along the eastern margin of the Heraklion Prefecture and in Messara graben to the south. Seismicity decreases rapidly from east to west. These observations are in agreement with the results presented by [13] and [21]. The intense seismic activity in the southern coastline of the Heraklion basin is probably associated with the offshore graben structure defined by E-W trending faults [23].

The overview of the near coast and onshore seismicity in Crete indicates a complex seismotectonic regime. This could be the result of both extensional and compressional stress fields in the area [22], resulting from the convergence of the European and African plates as well as from the extension within the Aegean basin.

## 6 Acknowledgments

This work was supported by the Greek General Secretariat of Research and Technology in the frame of Crete Regional Project 2000-2006 (M1.2): "An integrated system of seismic hazard monitoring and management in the front of the Hellenic arc", CRETE PEP\_7 (KP\_7). The GMT system [24] was used to plot part of the figures.

### References:

- [1] Papazachos, B.C., 1990, Seismicity of the Aegean and surrounding area, *Tectonophysics*, v. 178, pp. 287-308.
- [2] Papazachos, B.C., and Papazachou, C.B., 2003, *Earthquakes of Greece* (in Greek), Ziti, Thessaloniki, 273 p.
- [3] Papazachos, B.C., Karakostas, V.G., Papazachos, C.B., and Scordilis, E.M., 2000, The geometry of the Wadati-Benioff zone and lithospheric kinematics in the Hellenic arc, *Tectonophysics*, v. 319, pp. 275-300.
- [4] Faccenna, C., Jolivet, L., Piromallo, C., and Morelli, A., 2003, Subduction and the depth of convection of the Mediterranean mantle, *Journal of Geophysical Research*, v. 108, no. B2, p. 2099.
- [5] van Hinsbergen, D.J.J., Hafkenscheid, E., Spakman, W., Meulenamp, J.E., and Wortel, M.J.R., 2005, Nappe stacking resulting from subduction of oceanic and continental lithosphere below Greece, *Geology*, v. 33, pp. 325-328.
- [6] Le Pichon, X., and Angelier, J., 1979, The Hellenic Arc and trench system: a key to the neotectonic evolution of the Eastern Mediterranean area, *Tectonophysics*, v. 60, pp. 1-42.
- [7] Lyberis, N., and Lallemand, S., 1985, La transition subduction-collision le long de l'arc égéen externe, *Comptes Rendus de l'Académie des Sciences*, II, v. 300, pp. 887-890.
- [8] Finetti, I., 1985, Structure and evolution of the Central Mediterranean (Pelagian and Ionian Seas), in Stanley, D. J. and Wezel, F. C., eds., *Geological Evolution of the Mediterranean Basin*, Springer, pp. 215-230.
- [9] Sachpazi, M., Hirn, A., Clément, C., Haslinger, F., Laigle, M., Kissling, E., Charvis, P., Hello, Y., Lépine, J.-C., Sapin, M. and Ansorge, J. 2000. Western Hellenic subduction and Cephalonia Transform: local earthquakes and plate transport and strain, *Tectonophysics*, v. 319, pp. 301-319.

- [10] Mascle, J., Huguen, C., Benkhelil, J., Chomot-Rooke, N., Chaumillon, E., Foucher, J.P., Griboulard, R., Kopf, A., Lamarche, G., Volkonskaia, A., Woodside, J., and Zitter, T. 1999, Images may show start of European-African plate collision, *EOS Trans.*, v. 80, pp. 421-428.
- [11] Kokinou, E., and Kamberis, E., 2009, The structure of the Kythira-Antikythira strait, offshore SW Greece (35.7°-36.6N), in van Hinsbergen, D.J.J., Edwards, M.A., and Govers, R., eds., *Geodynamics of Collision and Collapse at the Africa-Arabia-Eurasia subduction zone*, *Geological Society of London Special Publication*, in press.
- [12] Papazachos, C.B. & Kiratzi, A.A. 1996, A detailed study of the active crustal deformation in the Aegean and surrounding area, *Tectonophysics*, v. 253, pp. 129-153.
- [13] Hatzfeld, D., Besnard, M., Makropoulos, K., and Hatzidimitriou, P., 1993, Microearthquake seismicity and fault-plane solutions in the southern Aegean and its geodynamic implications, *Geophysical Journal International*, v. 115, pp. 799-818.
- [14] Knapmeyer, M. and Harjes, H.P. (2000), Imaging crustal discontinuities and the downgoing slab beneath Crete, *Geophysical Journal International*, v. 143, pp. 1-21.
- [15] Makris, J., and Stobbe, C., 1984, Physical properties and state of the crust and upper mantle of the Eastern Mediterranean Sea deduced from geophysical data, *Marine Geology*, v. 55, pp. 347-363.
- [16] Jackson, J. (1994), Active tectonics of the Aegean region, *Annual Reviews of Earth and Planetary Science*, v. 22, pp. 239-271.
- [17] McClusky, S., Balassanian, S., Barka, A., Demir, C., Ergintav, S., Georgiev, I., Gurkan, O., Hamburger, M., Hurst, K., Kahle, H., Kastens, K., Kekeldze, G., King, R., Kotzev, V., Lenk, O., Mahmoud, S., Mishin, A., Nadariya, M., Ouzounis, A., Paradisses, D., Peter, Y., Prilepin, M., Reilinger, R., Sanli, I., Seeger, H., Tealeb, A., Toksoez, M.N., and Veis, G., 2000, Global Position System constraints on plate kinematics and dynamics in the eastern Mediterranean and Caucasus, *Journal of Geophysical Research*, v. 105, pp. 5695-5719.
- [18] Dewey, J.F., Helman, J.L., Turco, E., Hutton, D.W.H., and Knott, S.D., 1989, Kinematics of the Western Mediterranean, in Coward, M.P., and Dietrich, D., eds., *Alpine Tectonics*, *Geological Society, London, Special Publications*, v. 45, pp. 265-283.
- [19] ten Veen, J.H., and Meijer, P.Th., 1998, Late Miocene to Recent tectonic evolution of Crete (Greece): geological observations and model analysis, *Tectonophysics*, v. 298, pp. 191-208.
- [20] De Chabaliere, J.B., Lyon-Caen, H., Zollo, A., Deschamps, A., Bernard, P., and Hatzfeld, D., 1992, A detailed analysis of microearthquakes in western Crete from digital three-component seismograms, *Geophysical Journal International*, v. 110, pp. 347-360.
- [21] Delibasis, N., Ziazia, M., Voulgaris, N., Papadopoulos, T., Stavrakakis, G., Papanastassiou, D., Drakatos, G. (1999), Microseismic activity and seismotectonics of Heraklion area (Central Crete Island, Greece), *Tectonophysics*, v. 308(1-2), pp. 237-248.
- [22] Taymaz, T., Jackson, J. and Westway, R. (1990) Earthquake mechanisms in the Hellenic trench near Crete, *Geophysical Journal International*, v. 102, pp. 695-731.
- [23] Angelier, J., 1979, Neotectonique de l'arc Egeen, *Soc. Geol. du Nord*, 3.
- [24] Wessel, P., and Smith, W. H. F., 1998, New improved version of the Generic Mapping Tools Released, *EOS Trans. AGU*, 79, pp. 579.